

A STUDY ON INTEGER ADDITIVE SET-GRACEFUL GRAPHS

N. K. SUDEV

*Department of Mathematics
Vidya Academy of Science & Technology
Thalakkottukara, Thrissur, India.
E-mail: sudevnk@gmail.com*

K. A. GERMINA

*Department of Mathematics
University of Botswana
Gaborone, Botswana.
E-mail: srgerminaka@gmail.com*

Abstract

A set-labeling of a graph G is an injective set-valued function $f : V(G) \rightarrow \mathcal{P}(X)$, where X is a finite set and $\mathcal{P}(X)$ be its power set. A set-indexer of G is a set-labeling such that the induced function $f^\oplus : E(G) \rightarrow \mathcal{P}(X) - \{\emptyset\}$ defined by $f^\oplus(uv) = f(u) \oplus f(v)$ for every $uv \in E(G)$ is also injective. Let G be a graph and let X be a non-empty set. A set-indexer $f : V(G) \rightarrow \mathcal{P}(X)$ is called a set-graceful labeling of G if $f^\oplus(E(G)) = \mathcal{P}(X) - \{\emptyset\}$. A graph G which admits a set-graceful labeling is called a set-graceful graph. An integer additive set-labeling is an injective function $f : V(G) \rightarrow \mathcal{P}(\mathbb{N}_0)$, $\mathbb{N}_0$ is the set of all non-negative integers and an integer additive set-indexer is an integer additive set-labeling such that the induced function $f^+ : E(G) \rightarrow \mathcal{P}(\mathbb{N}_0)$ defined by $f^+(uv) = f(u) + f(v)$ is also injective. In this paper, we introduce the notion of integer additive set-graceful labeling of graphs analogous to the set-graceful labeling of graphs and study certain properties and characteristics of the graphs which satisfy this type of set-labeling.

Key words: Integer additive set-indexed graphs, set-graceful graphs, integer additive set-graceful graphs.

AMS Subject Classification : 05C78

1 Introduction

For all terms and definitions, not defined specifically in this paper, we refer to [7], [8], [14] and [20]. For more about graph labeling, we refer to [9]. Unless mentioned otherwise, all graphs considered here are simple, finite and have no isolated vertices.

The researches on graph labeling problems commenced with the introduction of β -valuations of graphs in [16]. Analogous to the number valuations of graphs, the concepts of set-labelings and set-indexers of graphs are introduced in [3] as follows.

Let X be a non-empty set and $\mathcal{P}(X)$ be its power set. A *set-labeling* of a graph G with respect to the set X is an injective set valued function $f : V(G) \rightarrow \mathcal{P}(X)$ with the induced edge function $f^+(E(G)) \rightarrow \mathcal{P}(X)$ is defined by $f^+(uv) = f(u) \oplus f(v)$ for all $uv \in E(G)$, where $\mathcal{P}(X)$ is the set of all subsets of X and $\oplus$ is the symmetric difference of sets. A set-labeling of G is said to be a *set-indexer* of G if the induced edge function f^+ is also injective. A graph which admits a set-labeling (or a set-indexer) is called a set-labeled (or set-indexed) graph. It is proved in [3] that every graph has a set-indexer.

A set-indexer $f : V(G) \rightarrow \mathcal{P}(X)$ of a given graph G is said to be a *set-graceful labeling* of G if $f^+(E(G)) = \mathcal{P}(X) - \{\emptyset\}$. A graph G which admits a set-graceful labeling is called a *set-graceful graph*.

The sumset of two non-empty sets A and B , denoted by $A+B$, is the set defined by $A+B = \{a+b : a \in A, b \in B\}$. If either A or B is countably infinite, then their sumset $A+B$ is also countably infinite. Hence all sets we consider here are non-empty finite sets. If $C = A+B$, where $A \neq \{0\}$ and $B \neq \{0\}$, then A and B are said to be the *non-trivial summands* of the set C and C is said to be a *non-trivial sumset* of A and B . Using the concepts of sumsets, the notion of integer additive set-labeling of a given graph G is defined as follows.

Definition 1.1. Let $\mathbb{N}_0$ be the set of all non-negative integers. Let $X \subseteq \mathbb{N}_0$ and $\mathcal{P}(X)$ be its power set. An *integer additive set-labeling* (IASL, in short) of a graph G is an injective function $f : V(G) \rightarrow \mathcal{P}(X)$ whose associated function $f^+ : E(G) \rightarrow \mathcal{P}(X)$ is defined by $f^+(uv) = f(u) + f(v)$, $uv \in E(G)$. A graph G which admits an integer additive set-labeling is called an integer additive set-labeled graph. (IASL-graph).

Definition 1.2. [12], [13] An *integer additive set-labeling* f is said to be an *integer additive set-indexer* (IASI, in short) if the induced edge function $f^+ : E(G) \rightarrow \mathcal{P}(X)$ defined by $f^+(uv) = f(u) + f(v)$ is also injective. A graph G which admits an integer additive set-indexer is called an *integer additive set-indexed graph*.

By an element of a graph G , we mean a vertex or an edge of G . The cardinality of the set-label of an element of a graph G is called the *set-indexing number* of that element. An IASL (or an IASI) is said to be a k -uniform IASL (or k -uniform IASI) if $|f^+(e)| = k \forall e \in E(G)$, where k is a positive integer. The vertex set $V(G)$ is called *l -uniformly set-indexed*, if all the vertices of G have the set-indexing number l .

With respect to an integer additive set-labeling (or integer additive set-indexer) of a graph G , the vertices of G has non-empty set-labels and the set-labels of every edge of G is the sumsets of the set-labels of its end vertices and hence no element of a given graph can have $\emptyset$ as its set-labeling. Hence, we need to consider only non-empty subsets of X for set-labeling the vertices or edges of G . Hence, all sets we mention in this paper are finite sets of non-negative integers. We denote the cardinality of a set A by $|A|$. We denote, by X , the finite ground set of non-negative

integers that is used for set-labeling the vertices or edges of G and cardinality of X by n .

In this paper, analogous to the set-graceful labeling of graphs, we introduce the notion of an integer additive set-graceful labeling of a given graph G and study its properties.

2 Integer Additive Set-Graceful Graphs

Certain studies have been done on set-graceful graphs in [3], [4], [5] and [6]. Motivated by these studies we introduce the notion of a graceful type of integer additive set-labeling as follows.

Definition 2.1. Let G be a graph and let X be a non-empty set of non-negative integers. An integer additive set-indexer $f : V(G) \rightarrow \mathcal{P}(X) - \{\emptyset\}$ is said to be an *integer additive set-graceful labeling* (IASGL) or a *graceful integer additive set-indexer* of G if $f^+(E(G)) = \mathcal{P}(X) - \{\emptyset, \{0\}\}$. A graph G which admits an integer additive set-graceful labeling is called an *integer additive set-graceful graph* (in short, IASG-graph).

As all graphs do not admit graceful IASIs in general, studies on the structural properties of IASG-graphs arouse much interest. The choice of set-labels of the vertices or edges of G and the corresponding ground set is very important to define a graceful IASI for a given graph.

A major property of integer additive set-graceful graphs is established as follows.

Property 2.2. *If $f : V(G) \rightarrow \mathcal{P}(X) - \{\emptyset\}$ is an integer additive set-graceful labeling on a given graph G , then $\{0\}$ must be a set-label of one vertex of G .*

Proof. If possible, let $\{0\}$ is not the set-label of a vertex in G . Since X is a non-empty subset of the set $\mathbb{N}_0$ of non-negative integers, it contains at least one element, say x , which is not the sum of any two elements in X . Hence, $\{x\}$ can not be the set-label of any edge of G . This is a contradiction to the hypothesis that f is an integer additive set-graceful labeling. $\square$

Examining the above mentioned property of IASG-graphs, it can be understood that if an IASL $f : V(G) \rightarrow \mathcal{P}(X) - \{\emptyset\}$ of a given graph G is an integer additive set-graceful labeling on G , then the ground set X must contain the element 0. That is, only those sets containing the element 0 can be chosen as the ground sets for defining a graceful IASI of a given graph.

Another trivial but an important property of certain set-labels of vertices of an IASG-graph G is as follows.

Property 2.3. *Let $f : V(G) \rightarrow \mathcal{P}(X) - \{\emptyset\}$ be an integer additive set-graceful labeling on a given graph G . Then, the vertices of G , whose set-labels are not the non-trivial sumsets of any two subsets of X , must be adjacent to the vertex v that has the set-label $\{0\}$.*

Proof. Let $A_i \neq \emptyset$ be a subset of X that is not a non-trivial sumset of any two subsets of X . But since f is an IASGL of G , A_i must be the set-label of some edge of G . This is possible only when A_i is the set-label of some vertex v_i that is adjacent to the vertex v whose set-label is $\{0\}$. $\square$

Invoking Property 2.3, we have the following remarks.

Remark 2.4. Let $x_i \in X$ be not the sum of any two elements in X . Since, f is an integer additive set-graceful labeling, $\{x_i\}$ must be the set-label of one edge, say e of G . This is possible only when one end vertex of e has the set-label $\{0\}$ and the other end vertex has the set-label $\{x_i\}$.

Remark 2.5. Let $f : V(G) \rightarrow \mathcal{P}(X) - \{\emptyset\}$ be an integer additive set-graceful labeling on a given graph G and let x_1 and x_2 be the minimal and second minimal non-zero elements of X . Then, by Remark 2.4, the vertices of G that have the set-labels $\{x_1\}$ and $\{x_2\}$, must be adjacent to the vertex v that has the set-label $\{0\}$.

Property 2.6. Let A_i and A_j are two distinct subsets of the ground set X and let x_i and x_j be the maximal elements of A_i and A_j respectively. Then, A_i and A_j are the set-labels of two adjacent vertices of an IASG-graph G is that $x_i + x_j \leq x_n$, the maximal element of X .

Proof. Let v be a vertex of G that has a set-label A_i whose maximal element x_i . If v is adjacent to a vertex, say u , with a set-label A_j whose maximal element is x_j , then $f^+(uv)$ contains the element $x_i + x_j$. Therefore, $x_i + x_j \in X$. Hence, $x_i + x_j \leq x_n$. $\square$

In view of Property, we can observe the following remarks.

Remark 2.7. Let $f : V(G) \rightarrow \mathcal{P}(X) - \{\emptyset\}$ be an integer additive set-graceful labeling on a given graph G and let x_n be the maximal element of X . If A_i and A_j are set-labels of two adjacent vertices, then $A_i + A_j$ is the set-label for the edge. Hence for any $x_i \in A_i, x_j \in A_j$, $x_i + x_j \leq x_n$ and hence, if one of the set is having the maximal element, then the other can not have a non-zero element. Hence, x_n is an element of the set-label of a vertex v of G if v is a pendant vertex that is adjacent to the vertex labeled by $\{0\}$. It can also be noted that if G is a graph without pendant vertices, then no vertex of G can have a set-label consisting of the maximal element of the ground set X .

The following results establish the relation between the size of an IASG-graph and the cardinality of its ground set.

Remark 2.8. Let f be an integer additive set-graceful labeling defined on G . Then, $f^+(E) = \mathcal{P}(X) - \{\emptyset, \{0\}\}$. Therefore, $|E(G)| = |\mathcal{P}(X)| - 2 = 2^{|X|} - 2 = 2(2^{|X|-1} - 1)$. That is, G has even number of edges.

Remark 2.9. Let G be an IASG-graph, with an integer additive set-graceful labeling f . By Remark 2.8, $|E(G)| = 2^{|X|} - 2$. Therefore, the cardinality of the ground set X is $|X| = \log_2[|E(G)| + 2]$.

The conditions for certain graphs and graph classes to admit a integer additive set-graceful labeling are established in following discussions.

Theorem 2.10. *A star graph $K_{1,m}$ admits an integer additive set-graceful labeling if and only if $m = 2^n - 2$ for any integer $n > 1$.*

Proof. Let v be the vertex of degree $d(v) > 1$. Let $m = 2^n - 2$ and $\{v_1, v_2, \dots, v_m\}$, be the vertices in $K_{1,m}$ which are adjacent to v . Let X be a set of non-negative integers containing 0.

First, assume that $K_{1,m}$ admits an integer additive set-graceful labeling, say f . Then, by Remark 2.8, $|E(G)| = m = 2^{|X|} - 2$. Therefore, $m = 2^n - 2$, where $n = |X| > 1$.

Conversely, assume that $m = 2^n - 2$ for some integer $n > 1$. Label the vertex v by the set $\{0\}$ and label the remaining m vertices of $K_{1,m}$ by the remaining m distinct non-empty subsets of X . Clearly, this labeling is an integer additive set-graceful labeling for $K_{1,m}$. This completes the proof. $\square$

The following theorem checks whether a tree can be an IASG-graph.

Proposition 2.11. *If a tree on m vertices admits an integer additive set-graceful labeling, then $1 + m = 2^n$, for some positive integer $n > 1$.*

Proof. Let G be a tree on m vertices. Then, $|E(G)| = m - 1$. Assume that G admits an integer additive set-graceful labeling, say f . Then, by Remark 2.8, for a ground set X of cardinality n , then $m - 1 = 2^{|X|} - 2$. Hence, $m + 1 = 2^{|X|}$. $\square$

Corollary 2.12. *Let G be a tree on m vertices. For a ground set X , let $f : V(G) \rightarrow \mathcal{P}(X)$ be an integer additive set-graceful labeling on G . Then, $|X| = \log_2(m + 1)$.*

Proof. Let G be a tree which admits an IASGL. Then, by Theorem 2.11, we have $m + 1 = 2^{|X|}$, where X is the ground set for labeling the vertices and edges of graphs. Hence, $|X| = \log_2(m + 1)$. $\square$

Theorem 2.13. *A tree G is an IASG-graph if and only if it is a star $K_{1,2^n-2}$, for some positive integer n .*

Proof. If $G = K_{1,2^n-2}$, then by Theorem 2.10, G admits an integer additive set-graceful labeling. Conversely, assume that the tree G on m vertices admits an integer additive set-graceful labeling, say f with respect to a ground set X of cardinality n . Therefore, all the $2^n - 1$ non-empty subsets of X are required for labeling the vertices of X . Also, note that $\{0\}$ can not be a set-label of any edge of G . Hence, all the remaining $2^n - 2$ non-empty subsets of X are required for the labeling the edges of G .

It is to be noted that the set-labels containing 0 are either the the sumsets of some other set-labels containing 0 or not a sumset of any subsets of X . Let $0 \in A_i \subseteq X$ be the set-label of an edge, say e , of G . Then, if A_i is not a sumset of subsets of X , by Property 2.3, then A_i must be the set-label of a vertex, say u , that is adjacent to the vertex v having set-label $\{0\}$. Assume that A_i is the sum set of two sets A_r and A_s . If $e = v_r v_s$, where A_r and A_s are respectively the set-labels of

v_r and v_s , then e will be an edge of G which is in a cycle of G , a contradiction to the fact that G is a tree. Therefore, vertices, whose set-labels containing 0, must be adjacent to the vertex v which has the set-label $\{0\}$.

Also, note that, by Remark 2.7, the vertices, whose set-labels containing the maximal element x_n of X must also be adjacent to the vertex labeled by $\{0\}$.

Let X_i be a subset of X which contains either 0 or x_n and let v_i be the vertex of G that has the set-label X_i . Then, the set-label of the edge vv_i is also X_i . Let X_j be a subset of X that contains neither 0 nor x_n and is the set-label of an edge e of G . Then, if X_j is not a sumset in $\mathcal{P}(X)$, then X_j must be the set-label of vertex which is adjacent to the vertex v having the set-label $\{0\}$. If X_j is the sumset of two subsets X_r and X_s and $e = v_r v_s$, where X_r and X_s are respectively the set-labels of v_r and v_s , then as explained above, the edge e will be in a cycle of G , a contradiction to the fact that G is a tree. Therefore, X_j must be the set-label of vertex which is adjacent to the vertex v having the set-label $\{0\}$.

Hence, all vertices of G having non-empty subsets of X , other than $\{0\}$, as the set-labels must be adjacent to the vertex v having set-label $\{0\}$. Hence, G is a star graph $\deg(v) = 2^n - 2$. $\square$

We now check the admissibility of integer additive set-graceful labeling by path graphs and cycle graphs.

Corollary 2.14. *For a positive integer $m > 2$, the path P_m does not admit an integer additive set-graceful labeling.*

Proof. Every path is a tree and no path other than P_2 is a star graph. Hence, by Theorem 2.13, P_m , $m > 2$ is not an IASG-graph. $\square$

Proposition 2.15. *For any positive integer $m > 3$, the cycle C_m does not admit an integer additive set-graceful labeling.*

Proof. Let X be a ground set with n elements. Since C_m has m edges, by Remark 2.8,

$$m = 2^n - 2 \tag{2.0.1}$$

Since C_m has no pendant vertices, by Proposition 2.7, the maximal element, say x_n , will not be an element of any set-label of the vertices of C_m . Therefore, only $2^{n-1} - 1$ non-empty subsets of X are available for labeling the vertices of C_m . Hence,

$$m \leq 2^{n-1} - 1 \tag{2.0.2}$$

Clearly, Equation 2.0.1 and Equation 2.0.2 do not hold simultaneously. Hence, C_m does not admit an integer additive set-graceful labeling. $\square$

An interesting question we need to address here is whether complete graphs admit integer additive set-graceful labeling. We investigate the conditions for a complete graph to admit an integer additive set-graceful labeling and based on these conditions check whether the complete graphs are IASG-graphs.

Theorem 2.16. *A complete graph K_m , does not admit an integer additive set-graceful labeling.*

Proof. Since K_2 has only one edge and K_3 has three edges, by Remark 2.8, K_2 and K_3 do not have an integer additive set-graceful labeling. Hence, we need to consider the complete graphs on more than three vertices.

Assume that a complete graph K_m , $m > 3$ admits an integer additive set-graceful labeling. Then, by Remark 2.8, $|E(G)| = 2^{|X|} - 2 = \frac{m(m-1)}{2}$. That is, $2^{|X|-1} - 1 = \frac{m(m-1)}{4}$. Since $|X| > 1$, $2^{|X|-1} - 1$ is a positive integer. Hence, $m(m-1)$ is a multiple of 4. This is possible only when either m or $(m-1)$ is a multiple of 4.

Since $|X| > 1$, $2^{|X|-1} - 1$ is a positive odd integer. Hence, for an odd integer k , either $m = 4k$ or $m-1 = 4k$. Therefore, $2^{|X|-1} - 1 = \frac{4k(4k-1)}{4} = k(4k-1)$ or $2^{|X|-1} - 1 = \frac{4k(4k-1)}{4} = k(4k+1)$. That is, $2^{|X|-1} = 1 + k(4k \pm 1)$. That is, if a complete graph K_m admits an integer additive set-graceful labeling, then there exist an integral solution for the equation

$$4k^2 \pm k + 1 = 2^n \quad (2.0.3)$$

where k is an odd non-negative integer and $n > 3$ be a positive integer.

The equation (2.0.3) can be written as a quadratic equation as follows.

$$4k^2 \pm k + (1 - 2^n) = 0 \quad (2.0.4)$$

The value of k is obtained from (2.0.4) as $k = \frac{\pm 1 \pm \sqrt{1-16(1-2^n)}}{8} = \frac{\pm 1 \pm \sqrt{2^n+4-15}}{8}$, which can not be a non-negative integer for the values $n > 3$. Hence, K_m does not admit an integer additive set-graceful labeling. $\square$

In this context, we need to find the conditions required for the given graphs to admit IASGLs. The structural properties of IASGL graphs are discussed in the following results.

Proposition 2.17. *Let G be an IASG-graph. Then, the minimum number of vertices of G that are adjacent to the vertex having the set-label $\{0\}$ is the number of subsets of X which are not the non-trivial sumsets of any subsets of X .*

Proof. Let G admits an integer additive set-graceful labeling f . Then, $f^+(E(G)) = \mathcal{P}(X) - \{\emptyset, \{0\}\}$. Let X_i be a non-empty subset of X , which is not a non-trivial sumset of any other subsets of X . Since G is an IASG-graph, X_i should be the set-label of some edge of G . Since X_i is the set-label of edges of G and is not a non-trivial sumset any two subsets of G , this is possible only when X_i is the set-label of some vertex of G which is adjacent to the vertex v whose set-label is $\{0\}$. Therefore, the minimum number of vertices adjacent to v is the number of subsets of X which are not the sumsets of any two subsets of X . $\square$

Another important structural property of an IASG-graph is established in the following theorem.

Theorem 2.18. *If a graph G admits an IASGL f with respect to a finite ground set X , then the vertices of G , which have the set-labels which are not non-trivial summands of any subset of X , are the pendant vertices of G .*

Proof. Let f be an IASGL defined on a given graph G . Then, every subset of X , other than $\emptyset$ and $\{0\}$, must be the set-label of some edges of G . Let X_i be not a non-trivial summand of any subset of X . Then, the vertex v_i with set-label X_i can not be adjacent to any other vertex v_j with set-label X_j , where $X_j \neq \{0\}$ as the set-label of the edge $v_i v_j$ is $X_i + X_j$ which is not a subset of X . Hence, v_i can be adjacent only to the vertex v having the set-label $\{0\}$. $\square$

Invoking the above theorems, we can establish the following result.

Theorem 2.19. *If G is an IASG-graph, then at least k pendant vertices must be adjacent to a single vertex of G , where k is the number of subsets of X which are neither the non-trivial sumsets of subsets of X nor the non-trivial summands of any subset of X .*

Proof. Let f be an IASGL defined on a given graph G . Then, every subset of X , other than $\emptyset$ and $\{0\}$, must be the set-label of some edges of G . If X_i is not a sumset of any two subsets of X , then by Proposition 2.17, the vertex v_i of G with the set-label X_i will be adjacent to the vertex v with the set-label $\{0\}$. If X_i is not a non-trivial summand of any subset of X , then by 2.18, the vertex v_i will be a pendant vertex. Therefore, the minimum number of pendant vertex required for a graph G to admit an IASGL is the number of subsets of X which are neither the non-trivial sumset of any two subsets of X nor the non-trivial summands of any subset of X . $\square$

The following result is an immediate consequence of the above theorems.

Theorem 2.20. *Let G be an IASG-graph which admits an IASGL f with respect to a finite non-empty set X . Then, G must have at least $|X| - 1$ pendant vertices.*

Proof. Let G admits an IASGL, say f with respect to a ground set $X = \{0, x_1, x_2, \dots, x_n\}$. By Theorem 2.19, the number of pendant vertices is equal to the number of subsets of X which are neither the non-trivial sumsets nor the non-trivial summands of any subsets of X . Clearly, the set $\{0, x_n\}$ is neither a sumset of any two subsets in X nor a non-trivial summand of any set in X . By Property 2.7, the vertex of G with set-label $\{0, x_n\}$ can be adjacent to a single vertex that has the set-label $\{0\}$. Now, consider the three element sets of the form $X_i = \{0, x_i, x_n\}$; $1 \leq i < n$. If possible let, the set X_i be the sumset of two subsets say A and B of X . Then, A and B can have at most two elements. Since $0 \in X_i$, 0 must belong to both A and B . Hence, let $A = \{0, a\}$ and $\{0, b\}$. Then, $X_i = A + B = \{0, a, b, a + b\}$ which is possible only when $a = b$ and hence $A = B$ and it contradicts the injectivity of f . Therefore, X_i is not a sumset of any other subsets of X . Since $x_n \in X_i$, by Property 2.7, it can not be a non-trivial summand of any subset of X . Therefore, X_i can be the set-label of a pendant vertex of G only. The number of three element subsets of X of this kind is $n - 1$. It is to be noted that be that a subset $\{0, x_i, x_j, x_n\}$ can be a sumset of two subsets $\{0, x_i\}$ and $\{0, x_j\}$ of X if $x_n = x_i + x_j$. This property holds for all subsets of X with cardinality greater than 3. Hence, the minimum number of subsets of X which are neither the non-trivial sumsets of any two subsets of X nor the non-trivial summands of any other subsets of X is $n = |X| - 1$. Hence, by Theorem 2.19, the minimum number of pendant vertices of an IASG-graph is $|X| - 1$. $\square$

An interesting question that arises in this context is about the existence of a graph corresponding to a given ground set X such that the function $f : V(G) \rightarrow \mathcal{P}(X)$ is a graceful IASI on G . Hence we introduce the following notion.

Definition 2.21. Let X be a non-empty finite set of non-negative integers. A graph G which admits a graceful IASI with respect to the set X is said to be a *graceful graph-realisation* of the set X with respect to the IASL f .

It can be noted that a star graph $K_{1,|X|-2}$ admits an IASGL. The question whether there is a non-bipartite graph that admits an IASGL with respect to a given ground set X is addressed in the following theorem.

Theorem 2.22. *Let X be a non-empty finite set of non-negative integers containing the element 0. Then, there exists a non-bipartite graceful graph-realisation G .*

Proof. Let X be a finite non-empty set of non-negative integers containing 0. Let $\mathcal{A}$ be the collection of all subsets of X which are not the non-trivial sumsets of any two subsets of X and let $\mathcal{B}$ be the collection of subsets of X which are not the non-trivial summands of any subsets of X . We need to construct an IASG-graph G with respect to X . For this, first take a vertex, say v_0 and label this vertex by $\{0\}$. Now, mark the vertices $v_1, v_2, \dots, v_r$, where $r = |\mathcal{A} \cup \mathcal{B}|$ and label these vertices in an injective manner by the sets in $\mathcal{A} \cup \mathcal{B}$. In view of Proposition 2.17 and Theorem 2.18, draw the edges from each of these vertices to the vertex v_0 . Next, mark new vertices $v_{r+1}, v_{r+2}, \dots, v_{r+l} = v_n$, where $l = |(\mathcal{A} \cup \mathcal{B})^c|$ and label these vertices in an injective manner by the sets in $(\mathcal{A} \cup \mathcal{B})^c$. Draw edges between the vertices v_i and v_j , if $f(v_i) + f(v_j) \subseteq X$, for all $0 \leq i, j \leq |V|$. Clearly, this labeling is an IASGL for the graph G constructed here. $\square$

Invoking all the above results, we can summarise a necessary and sufficient condition for a graph G to admit a graceful IASI with respect to a given ground set X .

Theorem 2.23. *Let X be a non-empty finite set of non-negative integers. Then, a graph G admits a graceful IASI if and only if the following conditions hold.*

- (a) $0 \in X$ and $\{0\}$ be a set-label of some vertex, say v , of G
- (b) the minimum number of pendant vertices in G is the number of subsets of X which are not the non-trivial summands of any subsets of X .
- (c) the minimum degree of the vertex v is equal to the number of subsets of X which are not the sumsets of any two subsets of X and not non-trivial summands of any other subsets of X .
- (d) the minimum number of pendant vertices that are adjacent to a given vertex of G is the number of subsets of X which are neither the non-trivial sumsets of any two subsets of X nor the non-trivial summands of any subsets of X .

Proof. The necessary part of the theorem follows together from Property 2.3, Proposition 2.17, Theorem 2.18 and 2.19 and the converse of the theorem follows from Theorem 2.22. $\square$

3 Conclusion

In this paper, we have discussed the concepts and properties of integer additive set-graceful graphs analogous to those of set-graceful graphs and have done a characterisation based on this labeling.

We note that the admissibility of integer additive set-indexers by the given graphs depends also upon the nature of elements in X . A graph may admit an IASGL for some ground sets and may not admit an IASGL for some other ground sets. Hence, choosing a ground set X is very important in the process of checking whether a given graph is an IASG-graph.

Certain problems in this area are still open. Some of the areas which seem to be promising for further studies are listed below.

Problem 3.1. Characterise different graph classes which admit integer additive set-graceful labelings.

Problem 3.2. Verify the existence of integer additive set-graceful labelings for different graph operations and graph products.

Problem 3.3. Analogous to set-sequential labelings, define integer additive set-sequential labelings of graphs and their properties.

Problem 3.4. Characterise different graph classes which admit integer additive set-sequential labelings.

Problem 3.5. Verify the existence of integer additive set-sequential labelings for different graph operations and graph products.

The integer additive set-indexers under which the vertices of a given graph are labeled by different standard sequences of non negative integers, are also worth studying. All these facts highlight a wide scope for further studies in this area.

References

- [1] K. Abhishek, *Set-Valued Graphs: A Survey*, Journal of Discrete Mathematical Sciences and Cryptography **18** (1-2)(2015), 55-80., DOI: 10.1080/09720529.2014.894306.
- [2] K. Abhishek, *Set-Valued Graphs II*, Journal of Fuzzy Set Valued Analysis **2013**(2013), 1-16., DOI: 10.5899/2013/jfsva-00149.
- [3] B. D. Acharya, **Set-Valuations and Their Applications**, MRI Lecture notes in Applied Mathematics, No.2, The Mehta Research Institute of Mathematics and Mathematical Physics, Allahabad, 1983.
- [4] B. D. Acharya, *Set-Indexers of a Graph and Set-Graceful Graphs*, Bull. Allahabad Math. Soc., **16**(2001), 1-23.

- [5] B. D. Acharya, K. A. Germina, K. L. Princy and S. B. Rao, *On Set-Valuations of Graphs*, In **Labeling of Discrete Structures and Applications**, (Eds.: B D Acharya, S Arumugam and A Rosa), Narosa Publishing House, New Delhi, 2008.
- [6] B. D. Acharya, K. A. Germina and K. Abhishek and P. J. Slater, *Some New Results on Set-Graceful and Set- Sequential Graphs*, Journal of Combinatorics, Information and System Sciences, **37**(2-4)(2012), 145-155.
- [7] J. A. Bondy and U. S. R. Murty, **Graph Theory**, Springer, 2008.
- [8] A. Brandstädt, V. B. Le and J. .P Spinrad, **Graph Classes:A Survey**, SIAM, Philadelphia, 1999.
- [9] J. A. Gallian, *A Dynamic Survey of Graph Labeling*, The Electronic Journal of Combinatorics (DS-16), 2013.
- [10] K. A. Germina and K. Abhishek, *Set-Valued Graphs I*, Journal of Fuzzy Set Valued Analysis, **2012**(2012), Article IDjfsva-00127, 17 pages.
- [11] K. A. Germina, K. Abhishek, K. L. Princy, *Further Results On Set-Valued Graphs*, Journal of Discrete Mathematical Sciences and Cryptography **11**(5)(2008), 559-566., DOI:10.1080/09720529.2008.10698208.
- [12] K. A. Germina and T. M. K. Anandavally, *Integer Additive Set-Indexers of a Graph:Sum Square Graphs*, Journal of Combinatorics, Information and System Sciences, **37**(2-4)(2012), 345-358.
- [13] K. A. Germina and N. K. Sudev, *On Weakly Uniform Integer Additive Set-Indexers of Graphs*, International Mathematical Forum, **8**(37)(2013) 1827-34.
- [14] F. Harary, **Graph Theory**, Addison-Wesley Publishing Company Inc., 1969.
- [15] S. M. Hegde, *On Set-Valuations of Graphs*, National Academy of Science Letters, **14**(4)(1991), 181-182.
- [16] A. Rosa, *On certain valuation of the vertices of a graph*, In **Theory of Graphs**, Gordon and Breach, Paris, 1967.
- [17] N. K. Sudev and K. A. Germina, *On Integer Additive Set-Indexers of Graphs*, International Journal Mathematical Sciences & Engineering Applications, **8**(2)(2014), 11-22.
- [18] N. K. Sudev and K. A. Germina, *Some New Results on Strong Integer Additive Set-Indexers*, Discrete Mathematics, Algorithms and Applications, **7**(1)(2015) 1-11., DOI: 10.1142/S1793830914500657.
- [19] N. K. Sudev, K. A. Germina and K. P. Chithra, *A Creative Review on Integer Additive Set-Valued Graphs*, International Journal of Scientific & Engineering Research, **6**(3)(2015), 372-378.
- [20] D. B. West, **Introduction to Graph Theory**, Pearson Education Inc., 2001.